

diabetic PAI-1  $+/+$  mice, is weak when compared with nondiabetic mice ( $\times 1.5$ ), and is hardly distinguishable from the induction observed in PAI-1  $-/-$  mice, at least for the eye (Fig. 3). Another striking result is that the level of TGF- $\beta$  expression and production by the kidney is higher in PAI-1  $-/-$  mice than in PAI-1  $+/+$  mice, in either basal or diabetic conditions. In the absence of pathologic data, including measurements of the glomerular basement membrane thickness, it is therefore difficult to conclude that PAI-1 knockout animals could be protected towards diabetic nephropathy and that this would be related to a lower concentration of TGF- $\beta$ .

ALEXANDRE HERTIG, GAELLE PELLE,  
and ERIC RONDEAU  
Paris, France

Correspondence to Dr. Alexandre Hertig, INSERM U702, Department of Nephrology A, Hôpital Tenon, 4, rue de la Chine, 75020 Paris, France.

E-mail: [alexandre.hertig@tnn.ap-hop-paris.fr](mailto:alexandre.hertig@tnn.ap-hop-paris.fr)

## REFERENCES

1. HERTIG A, RONDEAU E: Plasminogen activator inhibitor type 1: The two faces of the same coin. *Curr Opin Nephrol Hypertens* 13:39–44, 2004
2. NICHOLAS SB, AGUINIGA E, REN Y, *et al*: Plasminogen activator inhibitor-1 deficiency retards diabetic nephropathy. *Kidney Int* 67:1297–1307, 2005
3. LUTTUN A, LUPU F, STORKEBAUM E, *et al*: Lack of plasminogen activator inhibitor-1 promotes growth and abnormal matrix remodeling of advanced atherosclerotic plaques in apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol* 22:499–505, 2002
4. HERTIG A, BERROU J, ALLORY Y, *et al*: Type 1 plasminogen activator inhibitor deficiency aggravates the course of experimental glomerulonephritis through overactivation of transforming growth factor  $\beta$ . *FASEB J* 17:1904–1906, 2003

## Reply from the Authors

As we gain further insight into PAI-1-mediated mechanisms in disease, it is evident that PAI-1 may indeed have several functions. Early studies showed that PAI-1 deficiency protects against fibrosis, as in the bleomycin-induced pulmonary model [1], and this is supported by our studies in the streptozotocin-induced diabetic mouse model [2].

However, Hertig and Rondeau demonstrate that in inflammatory models of disease, PAI-1 deficiency may promote pathology. In fact, we have observed development of cardiac fibrosis in PAI-1  $-/-$  mice (unpublished data), as have Moriwaki *et al* [3], who also showed significant macrophage involvement in this process. These observations strongly suggest that PAI-1 function may be determined by the inciting process, and modulated by uPA expression, which may be tissue- or disease-specific.

The comment that induction of TGF- $\beta$  synthesis is weak ( $1.5 \times$ 's) reflects similar changes in total TGF- $\beta$  by Hertig *et al* ( $1.9 \times$ 's) [4]. Active TGF- $\beta$  was not mea-

sured in our paper. As such, it is provocative to assume that a seemingly “weak” stimulation of TGF- $\beta$  protein by PAI-1 may not have a significant clinical outcome (i.e., worsened albuminuria). Our data showed no significant difference in basal kidney TGF- $\beta$  expression, which was significantly reduced in diabetic PAI-1  $-/-$  mice (Fig. 3).

Significant structural changes of diabetic nephropathy, such as glomerular basement membrane thickness and fractional mesangial volume, are not typically observed after 4 weeks of diabetes in C57BL6 STZ-induced diabetic mice and, in fact, may not appear until  $\sim 6$  months of diabetes. For this reason, these parameters were not included in our study.

SUSANNE B. NICHOLAS and WILLA A. HSUEH  
Los Angeles, California

Correspondence to Susanne B. Nicholas, Warren Hall; David Geffen School of Medicine at UCLA, 900 Veteran Avenue, Suite 24-130, Los Angeles, CA 90095.

E-mail: [sunicholas@mednet.ucla.edu](mailto:sunicholas@mednet.ucla.edu)

## REFERENCES

1. EITZMAN D, MCCOY R, ZHENG X, *et al*: Bleomycin-induced pulmonary fibrosis in transgenic mice that either lack or overexpress the murine plasminogen activator-1 gene. *J Clin Invest* 97:232–237, 1996
2. NICHOLAS SB, AGUINIGA E, REN Y, *et al*: Plasminogen activator inhibitor-1 deficiency retards diabetic nephropathy. *Kidney Int* 67:1297–1307, 2005
3. MORIWAKI H, STEMPIEN-OTERO A, KREMEN M, *et al*: Overexpression of urokinase by macrophages or deficiency of plasminogen activator inhibitor type-1 causes cardiac fibrosis in mice. *Circ Res* 95:637–644, 2004
4. HERTIG A, BERROU J, ALLORY Y, *et al*: Type I plasminogen activator inhibitor deficiency aggravates the course of experimental glomerulonephritis through overactivation of transforming growth factor  $\beta$ . *FASEB J* 17:1904–1906, 2003

## Does fluvastatin really have an antioxidant effect in humans?

**To the Editor:** In the recent issue of *Kidney International*, Pat *et al* demonstrated that renal fibrosis after unilateral ureteral obstruction in rat was attenuated by the antioxidative effect of fluvastatin [1], a HMG-CoA reductase inhibitor notable for its additional antioxidant effect [2]. We undertook this clinical experiment because no in vivo human data are available addressing the antioxidant effect of fluvastatin. The patients studied consisted of 3 males and 3 females, with an average age of 56.5 years. All had biopsy-proven chronic glomerulonephritis with normal ranges of creatinine clearance and serum total protein. The mean urinary protein excretion was